

P C.1 (pg 2 of 6) Calorimetry

5. How many kilojoules of energy are required to change the temperature of 36 g of water from 5°C to 65°C?

Specific Heat Capacity (J/g°C)	
Liquid water	4.18
Liquid gasoline	2.22

- a. Compared to water, would more or less energy be required to change the temperature of 36 g of gasoline 60°C? Justify your response with a calculation.
6. If 125 g of water at 68°C is mixed with 138 g of gasoline at 17°C, set up an equation and then solve that equation to find the equilibrium temperature. (The specific heat capacity of gasoline is 2.22 J/g°C)
7. Equal masses of two different substances absorb the same amount of heat. The temperature of substance B increases twice as much as the temperature of substance A. Which substance has the higher specific heat capacity? EXPLAIN and/or SHOW CLEARLY.


8. Calculate the specific heat of a piece of metal with a mass of 125 g. The metal was heated in boiling water for an hour and was put into 250.0 ml of water that started at 20.5°C and reached equilibrium at 26.0°C. Fill in the temperature diagram to the right, show your problem set up and then solve. Circle your final answer.

Temperature Diagram	
T_M _____	
	ΔT_M _____
T_f _____	
	ΔT_w _____
T_w _____	

- a. From your list of specific heat capacities, select the two metals that are closest to the value you calculated. Propose which metal it is most likely to be the one tested in the procedure above. Justify your choice.

Specific Heat Capacity Values J/g°C	
sodium	1.22
aluminum	0.9
potassium	0.748
chromium	0.448
iron	0.447
nickel	0.443
copper	0.386
zinc	0.386
gallium	0.374
silver	0.236
cadmium	0.232
tin	0.22
barium	0.179
mercury	0.139
lead	0.138
platinum	0.134
gold	0.128

- b. After having made your choice in part a, and having been told that the students making the measurements did not stir their water metal mixture, is the arrangement of their thermometer most likely to be similar to diagram A or B. Justify your choice.



P C.1 (pg 3 of 6) Still More Calorimetry

Remember to get your work on paper before you even begin to touch a calculator.

Remember you may need to access reference pages for SHC or density values.

9. Determine the thermal equilibrium temperature (the specific heat capacity of gasoline is $2.22 \text{ J/g}^\circ\text{C}$)
 - a. If 100.0 g of water at 77°C is mixed with 80.0 g of water at 23°C
 - b. If 125 g of water at 68°C is mixed with 138 g of gasoline at 17°C , determine the thermal equilibrium temperature.
10. Determine the (Assume the density of alcohol = 2.7 g/ml)
 - a. volume of 11°C cold water that is combined with 145 ml of water at 78°C and thermal equilibrium is reached at 33°C .
 - b. size of the hot piece of aluminum If 99 ml of water at 21°C has a piece of hot aluminum at 106°C dropped into it, and the thermal equilibrium temperature is 29°C , determine the.
11. Determine the starting temperature (the specific heat capacity of alcohol = $2.46 \text{ J/g}^\circ\text{C}$)
 - a. of 205 g hot water, when mixed with 325 g of water at 13°C and the final temperature is 28°C .
 - b. of 100.0 g of alcohol, when mixed with 100.0 g of water at 45°C and the final temperature is 33°C .
12. How many joules of energy are required to change the temperature
 - a. of 1 g of water from 5°C to 10°C ?
 - b. of 23 g of water from 5°C to 6°C ?
13. What is the specific heat of (The specific heat capacity of olive oil is $2.17 \text{ J/g}^\circ\text{C}$)
 - a. olive oil if 435 J of heat is added to 3.4 g of 21.0°C olive oil that heats up to 85°C ?
 - b. some liquid when 785 J of heat is added to 13.4 g of this 21.0°C liquid that heats up to 75°C ?
14. How many kilojoules of heat is absorbed
 - a. when 1.00 L of water is heated from 18°C to 85°C ?
 - b. when 15.0 L of water is heated from 18°C to 55°C ?
15. What is the specific heat of
 - a. A piece of stainless steel with a mass of 1.55 g which absorbs 141 J of heat energy when its temperature increases by 178°C .
 - b. A piece of some metal with a mass of 8.67 g which absorbs 967 J of heat energy when its temperature increases by 157°C .
16. Determine the mass of a piece of lead if it drops from 100.0°C to 16.5°C in 120.0 g of water which started at 10.5°C . (The specific heat capacity of lead is $0.138 \text{ J/g}^\circ\text{C}$)
17. Determine the volume of water starting at 15.00°C that 200.0 g of aluminum starting at 100.0°C can heat to 27.0°C . (The specific heat capacity of aluminum is $0.900 \text{ J/g}^\circ\text{C}$)
18. What is the mass of a copper block if it was heated to 100.0°C and reached equilibrium with 200.0 ml of water at 33.5°C and the water started at 30.5°C . (The specific heat capacity of copper is $0.386 \text{ J/g}^\circ\text{C}$)
19. Determine the specific heat
 - a. for a 37.360 g piece of cold -16.0°C aluminum that cooled 155 ml of water from 63.0°C to 59.0°C .
 - b. Compare this experimental value to the theoretical value (listed on page 1) and determine the % error.
20. Calculate the specific heat of cadmium if a piece with a mass of 106 g heated to 100.0°C was put into 250.0 ml of water that started at 30.7°C and reached equilibrium at 32.0°C .
 - a. Compare this experimental value to the theoretical value and determine the % error.
 - b. Is it more likely that the error was caused by the Cd not being heated to 100.0°C , or was it heated higher than 100.0°C by accidentally passing it through the Bunsen flame before dropping it into the calorimeter?
21. A bowl of Rice Krispees provides 627 kJ of energy. What volume of water could the same amount of energy raise from 22.0°C to boiling temp?
22. A 65.2 g lump of aluminum that was put into 150.0 ml of water which rose from 12.3°C to 15.3°C . (The specific heat capacity of aluminum is $0.900 \text{ J/g}^\circ\text{C}$)
 - a. Determine the temperature change for this piece of metal
 - b. What temp did the aluminum start at?
23. A lump of aluminum (density = 2.7 g/ml) raised 150.0 ml of water from 23.0°C to 30.0°C . (The specific heat capacity of aluminum is $0.900 \text{ J/g}^\circ\text{C}$)
 - a. If the piece of aluminum displaces a volume of 23.2 ml of water, what was the temp change of the Al?
 - b. What was the starting temp of the Al?

In all of these problems, we must assume that the heat lost equals the heat gained.



–heat lost = heat gained

–(m × c × ΔT) = m × c × ΔT (Since the c is the same, all the problems are about water, you can cancel it out on both sides.)

1. For these problems, when the final equilibrium temp is not known you must give it the value of x. Remember that both the hot and the cold substance end at the same final temp which you do not know, so you must call it x. Using the x you can write the ΔT in terms of x. Refer to the temp diagram below. Since both substances are water, c has cancelled out.

a. $-(95)(x - 88) = (67)(x - 33)$
How is your algebra??? x = **65°C**



b. $-(125)(x - 68) = (138)(x - 17)$
How is your algebra??? x = **41°C**



2. In these problems you know the ΔT values, so you simply solve for the volume of the cold water. Since both substances are water, c has cancelled out.

a. $-(345)(43 - 78) = (x)(43 - 12)$
How is your algebra??? x = 390 g
which of course equals **390 ml** since the density of water is 1 g/ml



b. $-(99)(43 - 89) = (x)(43 - 31)$
How is your algebra??? x = 380 g
which of course equals **380 ml** since the density of water is 1 g/ml



3. In these problems, the algebra is a little bit faster if you set x equal to the ΔT, however this means you have one more step to finish the problem after solving for x. Since both substances are water, c has cancelled out.

a. $-(95)(x) = (128)(28 - 11)$
How is your algebra??? x = -23°C
but x was defined as the ΔT = T_f - T_i, thus -23 = 28 - T_f, thus T_f = **51°C**
OR you may have set it up this way $-(95)(28 - x) = (128)(28 - 11)$, and x = **51°**



b. $-(86)(68 - 98) = (56)(x)$
How is your algebra??? x = 46°C
but x was defined as the ΔT = T_f - T_i, thus 68 - T_i = 46°C, thus T_i = **22°C**
OR you may have set it up this way $-(86)(68 - 98) = (56)(68 - x)$ and x = **22°**



4. For these problems, when the final equilibrium temp is not known you must give it the value of x. Remember that both the hot and the cold substance end at the same final temp which you do not know, so you must call it x. Using the x you can write the ΔT in terms of x. Refer to the temp diagram below. In these problems the materials are different, so you must include the SHC values. Watch out for your T value which is quite different because the temp passes through 0°C. ΔT is the absolute value of the total number of degrees changed)

a. $-(95)(0.90)(x - 120) = (167)(4.18)(x - 33)$
How is your algebra??? x = **42.5°C**

b. $-(4.18)(55)(x - 65) = (0.138)(55)(x + 15)$
How is your algebra??? x = **62.4°C**



5. In this problem there is only one substance. This means you should use the heat formula $q = m \times c \times \Delta T$

$$\left(\frac{4.18J}{g^{\circ}C}\right)(36g)(65 - 5^{\circ}C) = 9028.8J \left(\frac{1kJ}{1000J}\right) = 9.03kJ$$

- a. The gasoline would require less energy because it has a smaller specific heat capacity and thus less energy is needed to move its temperature.

6. In this problem, there are both a hot and cold substances going through temperature changes. this requires using this equation: $-(m \times c \times \Delta T) = m \times c \times \Delta T$

$$-\left(\frac{4.18J}{g^{\circ}C}\right)(125g)(x - 68^{\circ}C) = \left(\frac{2.22J}{g^{\circ}C}\right)(138g)(x - 17^{\circ}C) \quad x = 49^{\circ}C$$

7. Substance A must have a higher specific heat capacity because an equal amount of energy causes it to change fewer degrees.

$$1g \times \#_A J \times 1^{\circ}C = 1g \times \#_B J \times 2^{\circ}C \quad \text{thus } A > B$$

Plug in some simple (absolute) values for ΔT , making ΔT of B twice as large as ΔT , and choose equal mass values.

If you wish, you can pick a simple value for the specific heat capacity of A, and solve for B

$$1 \times 1J/g^{\circ}C \times 1^{\circ} = 1 \times c_B \times 2^{\circ} \quad \text{so, } c_B = 0.5, \text{ thus } A > B$$

8. In this problem you can assume that boiling water is $100^{\circ}C$ and since the metal was in this water warming up for over an hour, the metal must also at $100^{\circ}C$. Solve this equation: $-(m \times c \times \Delta T) = m \times c \times \Delta T$ for c of the metal

$$c_m = -\left(\frac{(m_w)(c_w)(\Delta T_w)}{(m_M)(\Delta T_M)}\right) \quad \text{then substitute the values and solve } c_m = -\left(\frac{(250)(4.18)(26 - 20.5)}{(125)(26 - 100)}\right)$$

Solve for $c = 0.62 J/g^{\circ}C$

- a. Check out the values on the specific heat capacity list and the two closest values are potassium = $0.748 J/g^{\circ}C$ and chromium = $0.448 J/g^{\circ}C$ (or iron which is so close, $0.447 J/g^{\circ}C$). Given these two choices, the metal must be **chromium or iron**. Even though the experimental value is closer to the potassium value, it can not be potassium because if you placed potassium in water it would react and burst into flames, thus potassium would never be able to be heated up in boiling water.

- b. Since the experimental specific heat capacity is higher ($0.62 J/g^{\circ}C > 0.448 J/g^{\circ}C$) than the theoretical value, we need to select an error source that would have resulted in a higher calculation. If the thermometer were arranged as in diagram A, and the water were not stirred, the final temperature would be erroneously higher than if this error did not occur. An erroneously higher final temp would cause ΔT_M to be smaller and ΔT_w to be larger.

Transfer these problem into the formula and you will see that the erroneous experimental value turns up to be too large.

$$\left(\frac{(c_w)(m_w)(\Delta T_w)\uparrow}{(m_M)(\Delta T_M)\downarrow}\right) = c_m \quad \uparrow$$

Using the formula $Heat\ lost/gained = c \times M \times \Delta T$ if only one substance is involved,
or $-(c \times m \times \Delta T) = c \times m \times \Delta T =$ if there is both a hot and a cold substance involved.

- 9 a $-(4.18)(100)(x-77) = (4.18)(80)(x-23)$ $x = 53^{\circ}\text{C}$
b $-(4.18)(125)(x-68) = (2.22)(138)(x-17)$ $x = 49^{\circ}\text{C}$
- 10 a $-(4.18)(145)(33-78) = (4.18)(m)(33-11)$ $m = 297\text{ g}$ which is **297 ml** since the density of water = 1 g/ml
b $(4.18)(99)(29-21) = -(0.900)(m)(29-106)$ $m = 47.8\text{ g}$ and since $D = m/v$ so $m/D = V$ $47.8/2.7 = 17.6\text{ ml}$
- 11 a $(4.18)(325)(28-13) = -(4.18)(205)(\Delta T)$ $\Delta T = -23.8^{\circ}\text{C}$ but this is the ΔT , so $28 - (-23.8) = 51.8^{\circ}\text{C}$
OR you could put x in for the final temp: $(4.18)(325)(28-13) = (4.18)(205)(28-x)$ $x = 51.8^{\circ}\text{C}$
b $-(4.18)(100)(33-45) = (2.17)(100)(\Delta T)$ $\Delta T = 23^{\circ}\text{C}$ but this is the ΔT , so $33 - 23 = 10^{\circ}\text{C}$
OR you could put x in for the final temp: $-(4.18)(100)(33-45) = (2.46)(100)(33-x)$ $x = 12.6^{\circ}\text{C}$
- 12 a $(4.18)(1)(5) = 21\text{ J}$
b $(4.18)(23)(1) = 96\text{ J}$
- 13 a $(x)(3.4)(64) = 435$ $x = 2.0\text{ J/g}^{\circ}\text{C}$
b $(x)(13.4)(54) = 785$ $x = 1.08\text{ J/g}^{\circ}\text{C}$
- 14 a $(4.18)(1000)(67) = 280\text{ kJ}$
b $(4.18)(15,000)(37) = 2,319,900\text{ J} = 2300\text{ kJ}$
- 15 a $(c)(1.55)(178) = 141\text{ J}$ $c = 0.511\text{ J/g}^{\circ}\text{C}$
b $(c)(8.67)(157) = 0.710\text{ J/g}^{\circ}\text{C}$
- 16 $-(0.138)(m)(16.5-100) = (4.18)(120)(6)$ $m = 261\text{ g}$
- 17 $-(0.900)(200)(27-100) = (4.18)(m)(27-15)$ $m = 261.6\text{ g}$ and since the density of water is 1 g/ml $261.6\text{ g} = 262\text{ ml}$
- 18 $-(0.386)(m)(35.5-100) = (4.18)(200)(33.5-30.5)$ $m = 97.7\text{ g}$
- 19 (c) $(37.360)[59 - (-16)] = -(4.18)(155)(59-63)$ $c = 0.924\text{ J/g}^{\circ}\text{C}$
b $(0.924 - 0.900)/0.900 = \text{approx } 2.8\%$
- 20 a $-(c)(106)(32-100) = (4.18)(250)(32-30.7)$ $c = 0.188\text{ J/g}^{\circ}\text{C}$
b $(0.232 - 0.188)/0.232 = 19.0\%$
c Because the experimental c is smaller than the theoretical value, it is more likely that the error was caused by the Cd not being heated all the way through. This would cause the equilibrium temp to be too low which would cause the ΔT_m to be too large (which is on the bottom of the quotient) and the ΔT_w to be too small (which is on the top of the quotient), and all other values being the same (c_w and mass values) would result in a smaller SHC for cadmium.
- 21 $627\text{ kJ} \gg 627,000\text{ J} = (4.18)(m)(100-22)$ $m = 1920\text{ g}$ which is of course equal to **1920 ml**
- 22 a $(0.900)(65.2)(\Delta T) = (4.18)(150)(15.3-12.3)$ $\Delta T = -32.1^{\circ}\text{C}$
b Since the aluminum must have been hotter to cause the water to raise in temp, $15.3 - (-32.1) = 47.4^{\circ}\text{C}$
- 23 a Small density twist: for the aluminum, $D \times V = m$ $2.7\text{ g/ml} \times 23.2\text{ ml} = 62.64\text{ g}$
 $-(0.900)(62.64)(\Delta T) = (4.18)(150)(30-23)$ $\Delta T = -78.0^{\circ}\text{C}$
b Since the aluminum must have been hotter to cause the water's temp to raise, $30 - (-78) = 108.0^{\circ}\text{C}$